

1. Lyketsos CG, Lopez O, Jones B, Fitzpatrick AL, Breitner J, DeKosky S. Prevalence of neuropsychiatric symptoms in dementia and mild cognitive impairment: results from the cardiovascular health study. *JAMA* 2002;288(12):1475-83. Available from: <http://jamanetwork.com/journals/jama/fullarticle/195320>
2. van der Steen JT, Smaling HJA, van der Wouden JC, Bruinsma MS, Scholten RJPM, Vink AC. Music-based therapeutic interventions for people with dementia. *Cochrane Database of Systematic Reviews* 2018, Issue 7. Art. No.: CD003477. DOI: 10.1002/14651858.CD003477.pub4. <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD003477.pub4/full>
3. Karkou V, Meekums B. Dance movement therapy for dementia. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No.: CD011022. DOI: 10.1002/14651858.CD011022.pub2.
4. Deshmukh SR, Holmes J, Cardno A. Art therapy for people with dementia (Protocol). *Cochrane Database of Systematic Reviews* 2014, Issue 4. Art. No.: CD011073. DOI: 10.1002/14651858.CD011073.