

Stroke rehabilitation trials: let's measure what matters

Julie Duncan Millar for Evidently Cochrane 08 March 2018

References

Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J, van Wijck F. Interventions for improving upper limb function after stroke. *Cochrane Database of Systematic Reviews* 2014, Issue 11. Art. No.: CD010820. DOI: 10.1002/14651858.CD010820.pub2. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010820.pub2/full>

Santisteban L, Térémetz M, Bleton J, et al. Upper Limb Outcome Measures Used in Stroke Rehabilitation Studies: A Systematic Literature Review. *PloS one* 2016;11(5):e0154792.

Purton J. Stroke survivors' experiences of upper limb dysfunction: a longitudinal exploratory study, PhD Thesis, Keele University, UK, 2017.

(ICHOM –I've put the website in the blog) Salinas J, Sprinkhuizen S.M, Ackerson T, et al. 2016, An International Standard Set of Patient-Centered Outcome Measures After Stroke. *Stroke* 2016; 47(1); 180-186.

Kwakkel G, Lannin NA, Borschmann K, et al. Standardised measurement of sensorimotor recovery in stroke trials: consensus-based core recommendations from the Stroke Recovery and Rehabilitation Roundtable (SRRR) *International Journal of Stroke* 2017; 12(5); 451-461.