

## Retention to clinical trials: how can we keep participants involved?

Katie Gillies and Derek Stewart for Evidently Cochrane 18 May 2018

### References

Brueton VC, Tierney J, Stenning S, Harding S, Meredith S, Nazareth I, Rait G. Strategies to improve retention in randomised trials. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No.: MR000032. DOI: 10.1002/14651858.MR000032.pub2.

<http://cochranelibrary-wiley.com/doi/10.1002/14651858.MR000032.pub2/abstract>

Healy P, Galvin S, Williamson PR, Treweek S, Whiting C, Maeso B, Bray C, Brocklehurst P, Clarke Moloney M, Douiri A, Gamble C, Gardner HR, Mitchell D, Stewart D, Jordan J, O'Donnell M, Clarke M, Pavitt SH, Woodford Guegan E, Blatch-Jones A, Smith V, Reay H, Devane D. Identifying trial recruitment uncertainties using a James Lind Alliance Priority Setting Partnership – the PRioRiTty (Prioritising Recruitment in Randomised Trials) study. *Trials* 2018;19:147.

Available

from: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5831203/pdf/13063\\_2018\\_Article\\_2544.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5831203/pdf/13063_2018_Article_2544.pdf)

Tudur Smith C, Hickey H, Clarke M, Blazeby J, Williamson P. The trials methodological research agenda: results from a priority setting exercise. *Trials* 2014;15:32. Available

from: <https://link.springer.com/content/pdf/10.1186%2F1745-6215-15-32.pdf>