

## Muddled by mu-opioid antagonists for opioid-induced constipation? Me too...

Anna Sutherland for Evidently Cochrane - 17 August 2018

### References

[i] Candy B, Jones L, Vickerstaff V, Larkin PJ, Stone P. Mu-opioid antagonists for opioid-induced bowel dysfunction in people with cancer and people receiving palliative care. Cochrane Database of Systematic Reviews 2018, Issue 6. Art. No.: CD006332. DOI: 10.1002/14651858.CD006332.pub3.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006332.pub3/full>

[ii] National Institute for Health and Care Excellence. *Naldemedine for treating opioid-induced constipation ID1189*. London: National Institute for Health and Care Excellence; in development. (NICE Technology Appraisal Guidance in development – TA10291). <https://www.nice.org.uk/guidance/proposed/gid-ta10291> accessed 10/7/18

[iii] Janku F, et al, Treatment with methylnaltrexone is associated with increased survival in patients with advanced cancer, *Ann Oncol*. 2016 Nov;27(11):2032-2038

[iv] Miller, E. Easing the strain: put your feet up for constipation, 2016, Evidently Cochrane, <http://www.evidentlycochrane.net/feet-up-constipation/>. accessed 10/7/18