

## **Early additional food and fluids for healthy breastfed full-term infants: review and reflections**

**S. Chapman and A. Einion for Evidently Cochrane 04 November 2016**

### **References**

- De Jager E, Skouteris H, Broadbent J et al (2013). 'Psychosocial correlates of exclusive breastfeeding: a systematic review'. *Midwifery*, 29: 506-518.
- O'Brien M, Buikstra E and Hegney D (2008). 'The influence of psychological factors on breastfeeding duration'. *Journal of Advanced Nursing*, 63(4): 387-408.
- Smith HA and Becker GE (2016). 'Early additional food and fluids for healthy breastfed full-term infants: review'. *Cochrane Database of Systematic Reviews*, 8: DC006462.  
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006462.pub4/full>
- Wight N and Marinelli KA (2014). 'ABM Clinical Protocol #1: Guidelines for blood glucose monitoring and treatment of hypoglycemia in term and late-preterm neonates, revised 2014'. *Breastfeeding Medicine*, 9(4): 173-179.
- WHO (2011). *Exclusive breastfeeding for six months best for babies everywhere*, Geneva: WHO.  
[www.who.int/mediacentre/news/statements/2011/breastfeeding\\_20110115/en/](http://www.who.int/mediacentre/news/statements/2011/breastfeeding_20110115/en/)