Perineal pain after birth: what can help?

Jennifer Hanratty for Evidently Cochrane 9 May 2017

References updated 13 October 2020

Aasheim V, Nilsen ABV, Reinar LM, Lukasse M. Perineal techniques during the second stage of labour for reducing perineal trauma. Cochrane Database of Systematic Reviews 2017, Issue 6. Art. No.: CD006672. DOI: 10.1002/14651858.CD006672.pub3. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006672.pub3/full

Beckmann MM, Stock OM. Antenatal perineal massage for reducing perineal trauma. Cochrane Database of Systematic Reviews 2013, Issue 4. Art. No.: CD005123. DOI: 10.1002/14651858.CD005123.pub3. http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005123.pub3/abstract

Chou D, Abalos E, Gyte GML, Gülmezoglu AM. Paracetamol/acetaminophen (single administration) for perineal pain in the early postpartum period. Cochrane Database of Systematic Reviews 2013, Issue 1. Art. No.: CD008407. DOI: 10.1002/14651858.CD008407.pub2. http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008407.pub2/abstract

East CE, Dorward EDF, Whale RE, Liu J. Local cooling for relieving pain from perineal trauma sustained during childbirth. Cochrane Database of Systematic Reviews 2020, Issue 10. Art. No.: CD006304. DOI: 10.1002/14651858.CD006304.pub4. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006304.pub4/full

Gupta JK, Sood A, Hofmeyr GJ, Vogel JP. Position in the second stage of labour for women without epidural anaesthesia. Cochrane Database of Systematic Reviews 2017, Issue 5. Art. No.: CD002006. DOI: 10.1002/14651858.CD002006.pub4. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD002006.pub4/full

Jiang H, Qian X, Carroli G, Garner P. Selective versus routine use of episiotomy for vaginal birth. Cochrane Database of Systematic Reviews 2017, Issue 2. Art. No.: CD000081. DOI: 10.1002/14651858.CD000081.pub3. http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000081.pub3/abstract

McAndrew, F., Thompson, J., Fellows, L., Large, A., Speed, M., & Renfrew, M. J. (2012). Infant feeding survey 2010. *Leeds: Health and Social Care Information Centre*. http://content.digital.nhs.uk/catalogue/PUB08694/Infant-Feeding-Survey-2010-Consolidated-Report.pdf

Shepherd E, Grivell RM. Aspirin (single dose) for perineal pain in the early postpartum period. Cochrane Database of Systematic Reviews 2020, Issue 7. Art. No.: CD012129. DOI: 10.1002/14651858.CD012129.pub3. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012129.pub3/full

Wuytack F, Smith V, Cleary BJ. Oral non-steroidal anti-inflammatory drugs (single dose) for perineal pain in the early postpartum period. Cochrane Database of Systematic Reviews2016, Issue 7. Art. No.: CD011352. DOI: 10.1002/14651858.CD011352.pub2. http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD011352.pub2/abstract

UNICEF. Caring for your baby at night A guide for parents *NCT.org*: UNICEF UK Baby Friendly Initiative.