

Prelabour rupture of membranes: induce or wait?

Lisa Smith for Evidently Cochrane 11 May 2017

References

Campbell, D. (2017) Jeremy Hunt orders investigation into baby deaths at NHS trust. *The Guardian*, 12th April 2017. Available at: <https://www.theguardian.com/society/2017/apr/12/jeremy-hunt-orders-investigation-into-baby-deaths-at-nhs-trust> (accessed 19 April 2017).

Middleton, P., Shepherd, E., Flenady, V., McBain, R.D., Crowther, C.A. (2017) Planned early birth versus expectant management (waiting) for prelabour rupture of membranes at term (37 weeks or more). *Cochrane Database of Systematic Reviews* 2017, Issue 1. Art. No.: CD005302.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005302.pub3/full>

Plain language summary: http://www.cochrane.org/CD005302/PREG_it-better-baby-be-born-immediately-or-wait-labour-start-spontaneously-when-waters-break-or-after-37

National Institute for Health and Clinical Excellence (2014) *Intrapartum care. Clinical guideline 190*. London: NICE.

National Institute for Health and Clinical Excellence (2008) *Induction of labour. Clinical guideline 70*. London: NICE.

NHS England (2016) *Saving Babies' Lives: a care bundle for reducing stillbirth*. Leeds: NHS England.

RCOG (2012) *The Prevention of Early-onset Neonatal Group-B Streptococcal Disease*. Green-top Guideline 36. 2nd edition. Available at: https://www.rcog.org.uk/globalassets/documents/guidelines/gtg_36.pdf (Accessed 1st May 2017)