

## **Cardiopulmonary resuscitation - 30:2 or just keep going?** **Scott Munro for Evidently Cochrane 22 June 2017**

### **References**

Brouwer TF, Walker RG, Chapman FW, Koster RW. Association between chest compression interruptions and clinical outcomes of ventricular fibrillation out-of-hospital cardiac arrest. *Circulation* 2015;132:1030-1037

Berg RA, Sanders AB, Kern KB, et al. Adverse hemodynamic effects of interrupting chest compressions for rescue breathing during cardiopulmonary resuscitation for ventricular fibrillation cardiac arrest. *Circulation* 2001;104:2465-2470

Perkins GD, Lockey AS, de Belder MA, Moore F, Weissberg P, Gray H, for the Community Resuscitation Group. National initiatives to improve outcomes from out-of-hospital cardiac arrest in England. *Emerg Med J* 2016;33(7):448–51. doi: 10.1136/emered-2015-204847.

Nolan JP, Hazinski MF, Aickin R, Bhanji F, Billi JE, Callaway CW, et al. Part 1: Executive summary: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2015;95:e1–31. doi: 10.1016/j.resuscitation.2015.07.039.

Resuscitation Council (UK). Resuscitation Council Guidelines 2015. Available at <https://www.resus.org.uk/resuscitation-guidelines/> [accessed 6 June 2017].

Wissenberg M, Lippert FK, Folke F, et al. Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA* 2013;310:1377–84.

Zhan L, Yang LJ, Huang Y, He Q, Liu GJ. Continuous chest compression versus interrupted chest compression for cardiopulmonary resuscitation of non-asphyxial out-of-hospital cardiac arrest. Cochrane Database of Systematic Reviews 2017, Issue 3. Art. No.: CD010134. DOI: 10.1002/14651858.CD010134.pub2.  
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010134.pub2/full>