

Videolaryngoscopy: watch this space

Rufaro Ndokera for Evidently Cochrane blog 19 October 2017 (updated 27 September 2018)

References

Abdelgadir IS, Phillips RS, Singh D, Moncreiff MP, Lumsden JL. Videolaryngoscopy versus direct laryngoscopy for tracheal intubation in children (excluding neonates). Cochrane Database of Systematic Reviews 2017, Issue 5. Art. No.: CD011413. DOI: 10.1002/14651858.CD011413.pub2.
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD011413.pub2/full>

Lewis SR, Butler AR, Parker J, Cook TM, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation. Cochrane Database of Systematic Reviews 2016, Issue 11. Art. No.: CD011136. DOI: 10.1002/14651858.CD011136.pub2.
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD011136.pub2/full>

Lingappan K, Arnold JL, Fernandes CJ, Pammi M. **Videolaryngoscopy versus direct laryngoscopy for tracheal intubation in neonates.** Cochrane Database of Systematic Reviews 2018, Issue 6. Art. No.: CD009975. DOI: 10.1002/14651858.CD009975.pub3. <http://cochranelibrary-wiley.com/enhanced/exportCitation/doi/10.1002/14651858.CD009975.pub3>