

## Retention to clinical trials: how can we keep participants involved?

**Katie Gillies and Derek Stewart for Evidently Cochrane 18 May 2018 updated 1 April 2021**

### References

Gillies K, Kearney A, Keenan C, Treweek S, Hudson J, Brueton VC, Conway T, Hunter A, Murphy L, Carr PJ, Rait G, Manson P, Aceves-Martins M. Strategies to improve retention in randomised trials. *Cochrane Database of Systematic Reviews* 2021, Issue 3. Art. No.: MR000032. DOI: 10.1002/14651858.MR000032.pub3. Accessed 01 April 2021.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.MR000032.pub3/full>

Healy P, Galvin S, Williamson PR, Treweek S, Whiting C, Maeso B, Bray C, Brocklehurst P, Clarke Moloney M, Douiri A, Gamble C, Gardner HR, Mitchell D, Stewart D, Jordan J, O'Donnell M, Clarke M, Pavitt SH, Woodford Guegan E, Blatch-Jones A, Smith V, Reay H, Devane D. Identifying trial recruitment uncertainties using a James Lind Alliance Priority Setting Partnership – the PRioRiTy (Prioritising Recruitment in Randomised Trials) study. *Trials* 2018;19:147. Available

from: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5831203/pdf/13063\\_2018\\_Article\\_2544.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5831203/pdf/13063_2018_Article_2544.pdf)

Tudur Smith C, Hickey H, Clarke M, Blazeby J, Williamson P. The trials methodological research agenda: results from a priority setting exercise. *Trials* 2014;15:32. Available

from: <https://link.springer.com/content/pdf/10.1186%2F1745-6215-15-32.pdf>