

The third stage of labour: active or expectant management of care?

Linda Biesty for Evidently Cochrane August 2019

References

Begley CM, Gyte GML, Devane D, McGuire W, Weeks A, Biesty LM. Active versus expectant management for women in the third stage of labour. *Cochrane Database of Systematic Reviews* 2019, Issue 2. Art. No.: CD007412. DOI: 10.1002/14651858.CD007412.pub5.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD007412.pub5/full>

International Confederation of Midwives (ICM) and International Federation of Gynaecology and Obstetrics (FIGO). Prevention and treatment of post-partum haemorrhage: new advances for low resource settings - Joint statement. The Hague, the Netherlands: International Confederation of Midwives; 2006. Available from: <http://www.who.int/pmnch/events/2006/figo2006statementeng.pdf>

Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006 Apr 1;367(9516):1066-1074. Available from: [https://doi.org/10.1016/S0140-6736\(06\)68397-9](https://doi.org/10.1016/S0140-6736(06)68397-9)

Lazarus JV, Lalonde A. Reducing postpartum hemorrhage in Africa. *Int J Gynaecol Obstet* 2005 Jan;88(1):89-90. Available from: <https://doi.org/10.1016/j.ijgo.2004.09.016>.

National Institute for Health and Care Excellence. *Intrapartum care for healthy women and babies*. (NICE CG190) [Issued December 2014; last updated February 2017]. London: National Institute for Health and Care Excellence; 2014. Available from: <https://www.nice.org.uk/guidance/cg190>

World Health Organization. *Managing complications in pregnancy and childbirth: a guide for midwives and doctors*. 2nd edition. Geneva: World Health Organization; 2017. Available from: <http://www.who.int/iris/bitstream/10665/255760/1/9789241565493-eng.pdf?ua=1>