

## References

Department of Health, England. *Safer Maternity Care: the National Maternity Safety Strategy – progress and next steps*. London: Department of Health, England; 2017. Available from: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/662969/Safer\\_maternity\\_care\\_-\\_progress\\_and\\_next\\_steps.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/662969/Safer_maternity_care_-_progress_and_next_steps.pdf)

Fernandez Turienzo C, Bick D, Bollard M, Brigante L, Briley A, Coxon K, Cross P, Healey A, Mehta M, Melaugh A, Moulla J, Seed PT, Shennan AH, Singh C, Tribe RM, Sandall J. POPPIE: protocol for a randomised controlled pilot trial of continuity of midwifery care for women at increased risk of preterm birth. *Trials* 2019 May 14;20(1):271. doi: 10.1186/s13063-019-3352-1. Available from: <https://trialsjournal.biomedcentral.com/track/pdf/10.1186/s13063-019-3352-1>

Fernandez Turienzo C, Roe Y, Rayment-Jones H, Kennedy A, Forster D, Homer C, McLachlan H, Sandall J. Implementation of midwifery continuity of care models for Indigenous women in Australia: Perspectives and reflections for the United Kingdom. *Midwifery* 2019 Feb;69:110-112. doi: 10.1016/j.midw.2018.11.005.

Homer C, Brodie P, Sandall J, Leap N. *Midwifery Continuity of Care*. 2<sup>nd</sup> edition. Chatswood, NSW: Elsevier Australia; 2019.

Kildea S, Gao Y, Hickey S, Kruske S, Nelson C, Blackman R, Tracy S, Hurst C, Williamson D, Roe Y. Reducing preterm birth amongst Aboriginal and Torres Strait Islander babies: a prospective cohort study, Brisbane, Australia. *EclinicalMedicine* 2019 Jul 1;12:43-51. doi: <https://doi.org/10.1016/j.eclinm.2019.06.001>. Available from: <https://www.thelancet.com/action/showPdf?pii=S2589-5370%2819%2930094-X>

Kildea S, Simcock G, Liu A, Elgbeili G, Laplante DP, Kahler A, Austin MP, Tracy S, Kruske S, Tracy M, O'Hara MW, King S. Continuity of midwifery carer moderates the effects of prenatal maternal stress on postnatal maternal wellbeing: the Queensland flood study. *Arch Womens Ment Health* 2018 Apr;21(2):203-214. doi: 10.1007/s00737-017-0781-2.

McRae DN, Muhajarine N, Stoll K, Mayhew M, Vedam S, Mpofu D, Janssen PA. Is model of care associated with infant birth outcomes among vulnerable women? A scoping review of midwifery-led versus physician-led care. *SSM Popul Health* 2016 Mar 18;2:182-193. doi: 10.1016/j.ssmph.2016.01.007. Available from:

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5757823/pdf/main.pdf>.

Erratum in: *SSM Popul Health* 2017 Jul 17;3:817. Available from:

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6033258/pdf/main.pdf>

Medley N, Vogel JP, Care A, Alfirevic Z. Interventions during pregnancy to prevent preterm birth: an overview of Cochrane systematic reviews. *Cochrane Database of Systematic Reviews* 2018, Issue 11. Art. No.: CD012505. DOI: 10.1002/14651858.CD012505.pub2. Available from: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012505.pub2/full>

National Maternity Review, NHS England. *Better Births: improving outcomes of maternity services in England: a Five Year Forward View for maternity care*. London: National Maternity Review, NHS England; 2016. Available from: <https://www.england.nhs.uk/wp-content/uploads/2016/02/national-maternity-review-report.pdf>

NHS England. *Implementing Better Births: a resource pack for Local Maternity Systems*. London: NHS England, Five Year Forward View; March 2017. Available from: <https://www.england.nhs.uk/wp-content/uploads/2017/03/nhs-guidance-maternity-services-v1.pdf>

NHS England. *The NHS Long Term Plan*. London: NHS England; 2019. Available from: <https://www.longtermplan.nhs.uk/publication/nhs-long-term-plan/>

Rayment-Jones H, Murrells T, Sandall J. An investigation of the relationship between the caseload model of midwifery for socially disadvantaged women and childbirth outcomes using routine data - a retrospective, observational study. *Midwifery* 2015 Apr;31(4):409-17. doi: 10.1016/j.midw.2015.01.003.

Sandall J, Soltani H, Gates S, Shennan A, Devane D. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database of Systematic Reviews* 2016, Issue 4. Art. No.: CD004667. DOI: 10.1002/14651858.CD004667.pub5. Available from: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD004667.pub5/full>

WHO Reproductive Health Library. WHO recommendation on midwife-led continuity of care during pregnancy. Geneva, Switzerland: World Health Organization; 2016. Available from: <https://extranet.who.int/rhl/topics/improving-health-system-performance/implementation-strategies/who-recommendation-midwife-led-continuity-care-during-pregnancy>

World Health Organization. WHO recommendations: intrapartum care for a positive childbirth experience. Geneva, Switzerland: World Health Organization; 2018. Available from: <https://www.who.int/reproductivehealth/publications/intrapartum-care-guidelines/en/>

World Health Organization. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva, Switzerland: World Health Organization; 2016. Available from: [https://www.who.int/reproductivehealth/publications/maternal\\_perinatal\\_health/anc-positive-pregnancy-experience/en/](https://www.who.int/reproductivehealth/publications/maternal_perinatal_health/anc-positive-pregnancy-experience/en/)