

Preventing awareness during surgery: what's the latest evidence?

Jennifer Kielty for Evidently Cochrane 22 May 2020

References

Checketts MR, Alladi R, Ferguson K, Gemmell L, Handy JM, Klein AA, Love NJ, Misra U, Morris C, Nathanson MH, Rodney GE, Verma R, Pandit JJ; Association of Anaesthetists of Great Britain and Ireland. Recommendations for standards of monitoring during anaesthesia and recovery 2015: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia* 2016;71(1):85-93. doi:10.1111/anae.13316. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5063182/pdf/ANAE-71-85.pdf>

Lewis SR, Pritchard MW, Fawcett LJ, Punjasawadwong Y. Bispectral index for improving intraoperative awareness and early postoperative recovery in adults. Cochrane Database of Systematic Reviews 2019, Issue 9. Art. No.: CD003843. DOI: 10.1002/14651858.CD003843.pub4. <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD003843.pub4/full>

Pandit JJ, Andrade J, Bogod DG, Hitchman JM, Jonker WR, Lucas N, Mackay JH, Nimmo AF, O'Connor K, O'Sullivan EP, Paul RG, Palmer JH, Plaat F, Radcliffe JJ, Sury MR, Torevell HE, Wang M, Hainsworth J, Cook TM; Royal College of Anaesthetists; Association of Anaesthetists of Great Britain and Ireland. 5th National Audit Project (NAP5) on accidental awareness during general anaesthesia: summary of main findings and risk factors. *Br J Anaesth* 2014;113(4):549-59. doi: 10.1093/bja/aeu313. Available from: <https://bjanaesthesia.org/action/showPdf?pii=S0007-0912%2817%2930746-8>