

Quitting smoking: which types of behavioural support work best to help people stop?

Annika Theodoulou and Dr Jamie Hartmann Boyce for Evidently Cochrane 31 May 2021

References

1. World Health Organization. World Health Organization fact sheets: Tobacco. Geneva: World Health Organization; May 2020. Web. 07 May 2021
<http://www.who.int/news-room/fact-sheets/detail/tobacco>.
2. Cahill K, Lindson-Hawley N, Thomas KH, Fanshawe TR, Lancaster T. Nicotine receptor partial agonists for smoking cessation. Cochrane Database of Systematic Reviews 2016, Issue 5. Art. No.: CD006103. DOI: 10.1002/14651858.CD006103.pub7.
<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006103.pub7/full>
3. Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T. Nicotine replacement therapy versus control for smoking cessation. Cochrane Database of Systematic Reviews 2018, Issue 5. Art. No.: CD000146. DOI: 10.1002/14651858.CD000146.pub5.
<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD000146.pub5/full>
4. Hartmann-Boyce J, Livingstone-Banks J, Ordóñez-Mena JM, Fanshawe TR, Lindson N, Freeman SC, Sutton AJ, Theodoulou A, Aveyard P. Behavioural interventions for smoking cessation: an overview and network meta-analysis. Cochrane Database of Systematic Reviews 2021, Issue 1. Art. No.: CD013229. DOI: 10.1002/14651858.CD013229.pub2.
<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013229.pub2/full>